

*Moving Beyond Mindfulness: Defining
Equanimity as an Outcome Measure in
Meditation and Contemplative Research*


**Gaëlle Desbordes, Tim Gard, Elizabeth
A. Hoge, Britta K. Hölzel, Catherine
Kerr, Sara W. Lazar, Andrew Olendzki &
David R. Vago**

Mindfulness

ISSN 1868-8527

Mindfulness

DOI 10.1007/s12671-013-0269-8



Your article is protected by copyright and all rights are held exclusively by Springer Science +Business Media New York. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

Moving Beyond Mindfulness: Defining Equanimity as an Outcome Measure in Meditation and Contemplative Research

Gaëlle Desbordes · Tim Gard · Elizabeth A. Hoge · Britta K. Hölzel · Catherine Kerr · Sara W. Lazar · Andrew Olendzki · David R. Vago

© Springer Science+Business Media New York 2014

Abstract In light of a growing interest in contemplative practices such as meditation, the emerging field of contemplative science has been challenged to describe and objectively measure how these practices affect health and well-being. While “mindfulness” itself has been proposed as a measurable outcome of contemplative practices, this concept encompasses multiple components, some of which, as we review here, may be better characterized as equanimity. Equanimity can be defined as an even-minded mental state or dispositional tendency toward all experiences or objects, regardless of their

origin or their affective valence (pleasant, unpleasant, or neutral). In this article, we propose that equanimity be used as an outcome measure in contemplative research. We first define and discuss the inter-relationship between mindfulness and equanimity from the perspectives of both classical Buddhism and modern psychology and present existing meditation techniques for cultivating equanimity. We then review psychological, physiological, and neuroimaging methods that have been used to assess equanimity either directly or indirectly. In conclusion, we propose that equanimity captures potentially the most important psychological element in the improvement of well-being, and therefore should be a focus in future research studies.

G. Desbordes (✉)
Athinoula A. Martinos Center for Biomedical Imaging,
Department of Radiology, Massachusetts General Hospital,
149 Thirteenth Street, Boston, MA 02129, USA
e-mail: desbordes@gmail.com

T. Gard
Faculty of Psychology and Neuroscience, Maastricht University,
Maastricht, The Netherlands

E. A. Hoge · S. W. Lazar
Department of Psychiatry, Massachusetts General Hospital,
Boston, MA, USA

B. K. Hölzel
Institute for Medical Psychology, Charité - Universitätsmedizin,
Berlin, Germany

C. Kerr
Department of Family Medicine, Alpert Medical School,
Brown University, Providence, RI, USA

A. Olendzki
Barre Center for Buddhist Studies, Barre, MA, USA

D. R. Vago
Department of Psychiatry, Brigham & Women's Hospital,
Boston, MA, USA

Keywords Equanimity · Mindfulness · Emotion regulation · Meditation · Acceptance

Introduction

In recent years, there has been increasing interest in the concept of mindfulness as well as in meditation techniques which can be used to develop it. These practices are directly inspired from the Buddhist tradition. The definition of mindfulness has itself evolved over the course of history and across different Buddhist schools (Bodhi 2011; Carmody et al. 2009; Dreyfus 2011; Dunne 2011; Grabovac et al. 2011; Grossman 2008; Kang and Whittingham 2010; Olendzki 2011; Williams and Kabat-Zinn 2011). Various attempts have been made to describe what mindfulness is in modern psychological terms, the most frequently cited being from Jon Kabat-Zinn who defined mindfulness as “paying attention in a particular way: on purpose, in the present moment, and non-judgmentally” (Kabat-Zinn 1990). This definition has been frequently used

in mental health research. However, scientific and clinical interpretations of mindfulness are not fully aligned with those from the Buddhist tradition (Dreyfus 2011; Wallace 2006). Over the last 10 years, several components of the practice of mindfulness have been identified. For example, Bishop et al. (2004) proposed a two-component model of mindfulness, in which the first component is self-regulated attention, with the purpose of maintaining attention on present-moment experience (“paying attention in a particular way: on purpose, in the present moment”), and the second component is an attitude of openness and acceptance (“in a particular way... non-judgmentally”). Shapiro et al. (2006) proposed a three-component model which further dissociates attention from intention (“on purpose”), and also includes attitude (“in a particular way”) as a third component. Baer et al. (2008) proposed a five-facet model of mindfulness in which two of the facets are “nonjudging of inner experience” and “nonreactivity to inner experience.” These definitions all have in common a component corresponding to an “attitude of openness and acceptance.” In this paper, we propose that this specific component shares similarities with the classical Buddhist description of equanimity, and that it is an important construct for scientific investigation.

In this article, we will define and discuss the interrelationship between mindfulness and equanimity from the perspectives of both classical Buddhism and modern psychology. Our goal is to help clarify the understanding of the concept of equanimity and also to promote further discussion and exploration of this concept. The emerging field of contemplative science has been challenged to describe and measure the effects of meditation and other related contemplative practices and to explain their relevance for health and well-being. Even though these practices are of great interest to the scientific and medical communities and to society in general, objective measures to assess their beneficial outcomes are lacking. We believe that a clearer distinction between mindfulness and equanimity will not only aid researchers who use various methods to assess the impact of meditation-based interventions and the mechanisms through which they work, but also help clinicians better understand how to teach and use meditative techniques with patients.

Mindfulness and Equanimity in the Buddhist Tradition

In this section, we will review definitions of mindfulness and equanimity in the Buddhist tradition, provide an illustrative example of mindfulness and equanimity in everyday life, highlight important distinctions between equanimity and indifference, and describe some meditative and contemplative methods for cultivating equanimity drawn from the Buddhist traditions. The next section will focus on equanimity from the perspective of Western psychological theory.

Definitions of Mindfulness and Equanimity

The Buddhist tradition offers a very rich taxonomy of the different states, abilities, and qualities of the mind. According to early Buddhist texts such as the Abhidhamma, what is usually referred to, or experienced, as a unified mental experience is comprised of numerous factors (Bodhi 2000; Tsering 2006). One important factor—which may or may not be present at any given moment—is mindfulness (*smṛti* in Sanskrit, *sati* in Pali), which is *the quality of mind that one recollects continuously without forgetfulness or distraction while maintaining attention on a particular [mental] object* (Analayo 2003; Bodhi 2000; Buddhaghosa 2003; Dreyfus 2011; Nyanaponika Thera 1962; Wallace 2006, 2011). In other words, mindfulness corresponds to the higher-level awareness of present-moment sensory, affective, and cognitive experiences. Mindfulness can also be described as the ability or *process* of maintaining the object of attention in working memory, whether this object corresponds to an external sensory stimulus, an interoceptive sensation from inside the body, a chain of discursive thoughts going through the mind, or even a memory from the past (Dreyfus 2011).

In the Buddhist tradition, the term “equanimity” (*upeksha* in Sanskrit, *upekkha* in Pali) is a complex construct that has been given multiple definitions along the development of Buddhist thought. At its heart is the word for “eye” and “see,” with a prefix suggesting “gazing upon” or observing without interference. In the Theravada Buddhist literature (which will be our main focus in this paper due to the prevalence of this particular branch of Buddhism in current mindfulness-based clinical interventions), there are two main usages of the term “equanimity.” First, it can refer to “neutral feeling,” a mental experience that is neither pleasant nor unpleasant, and that involves neither intensifying nor dampening current mental states (Bodhi 2000, pp. 34, 116). This first usage of the term equanimity corresponds to the Western psychological notion of “neutral valence,” and is commonly experienced throughout any ordinary day. The second meaning of equanimity, the one of interest in this paper, corresponds to a mental state or trait that is not easily achieved and typically requires some form of practice. It is “a state of mind that cannot be swayed by biases and preferences” (Bodhi 2000, p. 34), an “even-mindedness in the face of every sort of experience, regardless of whether pleasure [or] pain are present or not” (Thanissaro Bhikkhu 1996, p. 262). This state of equanimity manifests as “a balanced reaction to joy and misery, which protects one from emotional agitation” (Bodhi 2005, p. 154).

In brief, to rephrase this definition in modern terms, *equanimity is an even-minded mental state or dispositional tendency toward all experiences or objects, regardless of their affective valence (pleasant, unpleasant or neutral) or source*. Here, we use the term even-mindedness in its common definition as a state of being calm, stable and composed. Equanimity also involves a level of *impartiality* (i.e., being not

partial or biased), such that one can experience unpleasant thoughts or emotions without repressing, denying, judging, or having aversion for them. Similarly, in a state of equanimity, one can have pleasant or rewarding experiences without becoming over-excited (e.g., to the point of mania or hypomania), or trying to prolong these experiences, or becoming addicted to them. Grabovac et al. (2011) describe equanimity as “approaching pleasant, unpleasant and neutral experiences with equal interest.” It should be noted that this goes against our habitual tendencies to seek the pleasant and stay away from the unpleasant. In the Buddhist perspective, both of these tendencies are forms of craving (*tanha* in Pali, literally “thirst”), which constitutes the origin of all forms of suffering and dissatisfaction (*dukkha*) (Bodhi 2000, p. 289; Analayo 2003, chapter 13). In addition, it should be noted that the ideal form of equanimity embraced by Buddhism also includes having an equal attitude towards all beings, without the boundaries that we habitually draw between friends, strangers, and those we consider “difficult people,” in other words, “regarding all beings as equal in their right to have happiness and avoid suffering” (Tsering 2006, p. 86) and “treating them free from discrimination, without preferences and prejudices” (Bodhi 2000, p. 87). This aspect of equanimity can be cultivated with specific contemplative practice methods (see “Methods from the Buddhist Traditions for Cultivating Equanimity”).

We would like to emphasize that our proposed definition of equanimity as an even-minded state is both (1) a mental attitude of openness, even-mindedness, and acceptance that one purposefully cultivates (e.g., during formal meditation practice and/or throughout daily activities) and (2) an enduring state or trait that is the end result of this form of training. Indeed, the Buddhist tradition proposes that with practice, equanimity (along with the other “boundless qualities” of love, compassion, and sympathetic joy) becomes effortless and need no longer be purposefully invoked (Thrang Rinpoche 2002).

Illustrative Example of Mindfulness and Equanimity

To illustrate these concepts of mindfulness and equanimity with a simple example from everyday life, let us consider what happens when we are driving a car. It is possible to get in the car and arrive at our destination, lost in thought the whole time and completely unaware of what our hands and feet were doing or of what we saw or heard along the way. Through learning and habit, we keep the car on the road and respond to other vehicles, but conscious awareness is otherwise engaged. Such an extreme (but not uncommon) case of mind wandering corresponds to the absence of mindfulness. Alternatively, if we are mindful while driving, we can direct our attention to what is happening as we drive—our posture, the speed of the car, the cars around us, pedestrians, etc. We might also be aware of thoughts or emotions that may pass through our mind without getting lost in them. As we drive, we can choose to

actively note these events as they pass through our field of awareness. In this case, we are experiencing the flow of sensory and mental objects with an impartial yet caring attitude (equanimity) and mindful awareness.

If another driver suddenly cuts in front of us in a dangerous way, we might get scared or angry and yell or honk our horn as we hit the brakes, in which case our mindfulness and equanimity are disrupted by the intrusion of emotions such as anger or fear. Without mindfulness, we do not realize that these emotions are affecting our behavior and mood, causing perhaps road rage or panic. With mindfulness, we might gain awareness that our emotional state has changed, that our voice has gotten louder, that our muscles are more contracted, and that negative thoughts about the other driver are now passing through our mind. Being thus mindful of the arising of anger, we may (or may not) be able to maintain a state of equanimity and watch the experience of anger unfold and pass on its own accord. In the absence of equanimity, the experience of anger may become overwhelming or persist for a significant amount of time after the situation is over.

It is important to note that Buddhist psychology clearly distinguishes between mindfulness and equanimity. Each brings a different quality of mind to bear upon experience. Mindfulness emphasizes the ability to remain *consciously aware* of what is happening in the field of experience, while equanimity allows awareness to be *even and unbiased* by facilitating an attitude of non-attachment and nonresistance.

For the beginning meditator confronted with an emotionally charged situation, a nonjudging equanimous attitude is fleeting or often absent, even when mindfulness is present. This stage of the training may become discouraging to novices as they become increasingly aware of their disruptive thought patterns and impulsive—often maladaptive—actions, but do not yet have the equanimity to simply let the emotions pass through without reacting to them. A beginning meditator may then turn to other forms of regulation by actively suppressing or avoiding the emotions. An understanding of mindfulness and equanimity as separate skills that may develop at different rates might therefore be helpful to novices and teachers: the cultivation of mindfulness can be seen as the foundation on which equanimity will gradually develop.

Equanimity Versus Indifference in Buddhist Psychology

Buddhist psychology draws clear distinctions between equanimity and indifference. We review these distinctions in this section, and will discuss further below related views from within the framework of modern Western psychology.

Equanimity is sometimes described in the Buddhist literature as a *perspective shift* or a *sense of [temporal] detachment from the ongoing experience* (Bodhi 2000; Gunaratana 2002; Wallace 2006). Because of some connotations of the word “detachment” in English, which can imply “indifference,” this

has generated confusion. In a common misunderstanding of contemplative practices, equanimity is assumed to imply the absence of emotional reactivity, if not the absence of emotions themselves. In the above quote, “detachment” refers instead to elimination of craving or clinging to the objects of desire or discontent. Buddhist texts clearly distinguish between equanimity and indifference, where the latter implies a certain degree of apathy or lethargy. From the perspective of Buddhist philosophy, indifference is an “unwholesome” (i.e., detrimental or harmful) mental state which is due to ignorance of the true nature of things (Bodhi 2000). Indifference is explicitly warned against as the “near-enemy” of equanimity, which indicates that these two notions can seem deceptively similar at first but should be considered very different (Bodhi 2000, p. 87). The notion that equanimity includes a sense of care and attentiveness rather than indifference is illustrated in the following metaphor from the classical Buddhist tradition. Loving-kindness (*metta*) is likened to the sentiment felt by a mother toward a newborn infant (May she be well! May he thrive!); compassion (*karuna*) is the feeling extended to a sick child (May she be free from pain and suffering!); sympathetic joy (*mudita*) is how a mother feels toward a grown boy who leaves home to marry (Though it may be painful for me, I feel joy for him!); and equanimity is how a mother might feel on hearing about her grown child’s business dealings—she is attentive and caring about his welfare (not disengaged or indifferent), yet has no emotional entanglement to the content of the news she hears (Nanamoli 1991, p. 314). Therefore, equanimity is “not an attitude of cold indifference, but rather of mental imperturbability” (Thanissaro Bhikkhu 1996, p. 263). Indifference may even be considered a subtle form of aversion (Salzberg 1995, p. 150; Gunaratana 2002, p. 128).

Importantly, having equanimity does not mean suppressing emotions or giving up the affective coloring of our life experiences. When we are driving, for example, experiencing fear as another driver dangerously cuts in front of us is a more adaptive response than remaining indifferent to it. This fear is useful for stimulating a quick reaction to hit the brakes and avoid collision. However, once the danger has passed, it is more adaptive to bring attention back to current experience rather than continue to dwell in disruptive emotions. With equanimity, the transgression of the other driver is noted, but without the automatic seasonings of judgment, blame, resentment, and anger, much of which is driven by taking the incident as a personal insult. An even more skillful response, Buddhists would say, might be to forgive the person, have compassion for whatever conditions are compelling them to act so rudely, and even wish them well! Taking another example, being indifferent to the loss of a loved one could be considered pathological, because the experience of grief, although a disruptive and difficult emotion, is a healthy response of adjustment—as long as it does not persist indefinitely.

In the Buddhist view, the fundamental distinction between equanimity and indifference is also important in the context of equanimity towards people. Remaining indifferent to other people’s needs or their suffering is certainly not considered a form of equanimity, and is contrary to the Buddhist teachings. Buddhist masters recommend practicing equanimity towards beings by contemplating how all beings are similar to ourselves in that they, too, wish to gain happiness and be free of suffering, regardless of whether we would consider them our friends, enemies, or strangers (e.g., Dalai Lama 2001a, b). It follows that all beings should be treated with a very deep sense of care. Here we can see the link between equanimity, loving-kindness, and compassion. For more on the Buddhist practices for cultivating equanimity, see “[Methods from the Buddhist Traditions for Cultivating Equanimity](#).”

It is important to recognize the distinction between the affective tone of pleasure or pain, and one’s emotional response to it. There are many situations in which it is entirely appropriate to feel joyful, and we naturally welcome this experience because it involves pleasure. It can also be appropriate to feel sad, such as when grieving the loss of a loved one. From a Buddhist perspective, what determines our degree of “suffering” or dissatisfaction (*dukkha*) is our emotional response to joyfulness and sadness. For example, if we crave for the continuation or intensification of a joyful experience, then when it inevitably changes we will experience dissatisfaction in direct proportion to the strength of our craving. Similarly, psychological suffering occurs when an experience of sadness, for example, is amplified by aversion, resistance, denial, and other forms of unhealthy or unskillful emotional responses. Just as stress is defined not by what is happening in the world but by one’s response to what is happening (Lazarus 1991), so also suffering results not from the content of experience but from the emotional disposition with which it is experienced. Equanimity enables a skillful emotional response to the full range of feeling tones. Feeling happy without interfering with the mental pleasure by trying to grasp hold of it, or feeling unhappy and being able to “gaze upon” it with equanimity rather than trying to change it into something other than it is, are both skills that can be developed. Mindfulness is the first stage of this learning process by means of which one is able to focus attention upon the flow of experience without distraction. With practice, one will develop a greater ability to respond to this flow with equanimity.

Methods from the Buddhist Traditions for Cultivating Equanimity

The even mindedness that characterizes equanimity can be cultivated in various ways. In its most basic and accessible form, equanimity can be practiced in any situation when one is mindful of an object without being attracted to or repelled by it. One notices its presence without an intentional stance of

either favoring or opposing it, in other words, nonjudgmentally. Further development of this skill can proceed by either concentration (*samatha*) or insight (*vipassana*) practices, and the early Buddhist texts describe both paths in some detail (Buddhaghosa 2003; Wallace 2010). For example, in Theravada Buddhism, Vipassana meditation leads the practitioner through progressive stages of understanding the conditioned nature of all experience, culminating in an abiding and sustainable equanimity toward all phenomena which breaks the bonds of craving (for what one likes to continue or for the ending of what one does not like) that cause suffering. The path of concentration (*shamatha*) proceeds through four successive stages of intensifying meditative states, wherein a practitioner uses one-pointed focused concentration as a means to train the mind gradually from ordinary conceptual modes of operation to greater stillness and equanimity. According to the classical Buddhist literature, when one can maintain single-pointed concentration in the absence of mind wandering for some amount of time, one will first experience a great physical and mental pleasure that naturally arises as the mind suspends the construction of negative emotions; next, the rapture of this mental pleasure deepens as the mind lets go of conceptual thinking entirely and becomes even more absorbed in single-pointed concentration; then, the exuberance of this pleasure matures and mellows into a subtle but profound sense of happiness; while in the fourth and final stage, all pleasure is abandoned and an abiding mental equanimity is established. It is a process analogous to the purification of gold by burning off its contaminants, and a mind thus “purified” is said to be simultaneously bright, calm, steady, and flexible (Buddhaghosa 2003; Wallace 2010).

The advanced meditative states described above are usually achieved by practicing so-called nonreferential or nonconceptual forms of meditation, which have also been referred to as open presence, open awareness, nondual awareness, or nondual experience (Austin 2009; Dunne 2011; Josipovic 2010; Lutz, Dunne, and Davidson 2007). These practices are central to several branches of Buddhism such as Theravada, Zen, Mahamudra, and Dzogchen, and somewhat similar practices can also be found in non-Buddhist traditions such as the Vedic tradition and Chinese Qigong (Travis and Shear 2010). These meditative practices are said to make the mind more sensitive and flexible, which in turn facilitates the cultivation of beneficial states and traits such that the meditator becomes less easily disturbed by emotional states (Lutz et al. 2007). In the words of the Dalai Lama:

[The experience of Dzogchen] will give you the ability to prevent yourself from being overwhelmed by circumstances, good or bad. You will not fall into extreme states of mind: you will not get over-excited or depressed. Your attitude toward circumstances and events

will be as if you were someone observing the mind, without being drawn away by circumstances... You are not preoccupied by what arises in the mind, nor does it cause you any distress. You are free from conceptuality or any form of objectifying. And so it really does help you, in allowing you to be free from being caught up in the play of emotions (Dalai Lama 2001c, p. 162).

The interested reader is referred to the extensive literature on this topic from the contemplative traditions, which is beyond the scope of the present review.

In other Buddhist texts, including those from the Mahayana traditions, more accessible forms of equanimity practice are introduced in the early stages of meditation training. These methods are analytical (or conceptual) contemplations, also called “relative” practices, and can be used to complement the more advanced nonconceptual meditative practices mentioned above (Makransky 2007). Some of these contemplations are described in detail in the classic *Intermediate Stages of Meditation* by Kamalashila, an Indian master who wrote several treatises on introductory Buddhist practices (Dalai Lama 2001a; Thrangu Rinpoche 2002). One of the techniques for developing equanimity consists in trying to develop an impartial attitude toward all beings by reflecting on the “very simple truth” that “all of us, including the most despicable people, do the things we do because we’re seeking happiness and want to be free of suffering” (Wallace 2010, p. 155). A more detailed contemplation would be: “All beings are the same as ourselves in three basic ways. (1) They all want to be well, happy, and free of suffering. (2) They are all lost in self-centered reactions to their own narrow thoughts of self and others. (3) They all possess an innate capacity of goodness that has been obscured by the self-centered reactions” (Makransky 2007, p. 113; see also Dalai Lama 2001a, b).

Another classic analytical contemplation for developing equanimity presented by Kamalashila consists in reflecting upon the mind’s automatic classification of other people as “friends,” “enemies,” and “strangers,” and realizing that these categories are rather arbitrary and subject to change following life circumstances. In order to do so, it is considered effective to meditate on three particular individuals personally known by the meditator: one who has done him harm (“enemy” or “difficult person”), one who has benefited him directly (“friend”), and one neutral person (“stranger”). Observing how the mind reacts to thinking about each, the practitioner notices that, upon thinking of the enemy, the mind becomes irritated, resentful, or hateful; upon thinking of the friend, the mind becomes relaxed, comfortable, delighted; upon thinking of the stranger, there is neither irritation nor delight. The next step is to look for the reasons the mind might have for these types of responses. The practitioner will find that “these reasons are in fact superficial and based on narrow, self-serving attitudes.” At this point, “when we view things from a broader perspective with more

farsightedness, equanimity will dawn in our minds, enabling us to see the futility of hostility and clinging desire” and to “equalize our feelings toward those three individuals.” The next step in this technique is to “gradually extend the scope of this meditation” to our neighbors, our fellow citizens, and eventually “all the beings in the world” (Dalai Lama 2001a, p. 53; see also Wallace 2010, pp. 157–161, and Makransky 2007, p. 95–130). We should note that this practice is not unique to the Tibetan tradition, as it can also be found in Theravadan texts such as (Buddhaghosa 2003, p. 310).

Equanimity in Western Psychology and Psychophysiology

Equanimity has not been discussed much in the context of Western psychological theory. However, it appears in related concepts such as acceptance, nonjudgment, nonstriving, and nonreactivity—especially in the context of mindfulness meditation practice. Equanimity has also been conflated with emotional control and emotional suppression, which understandably can create confusion. Here, we attempt to clarify the concept of equanimity using the (Theravadan) Buddhist framework, but in the context of modern psychology and neuroscientific processes. We specifically propose and discuss the following: (1) equanimity is associated with a critical change in relation to one’s perceived experience; (2) equanimity shares some common effects with emotion regulation strategies; and (3) equanimity affects the time course of emotional and physiological response to a stressor.

Equanimity: A Relation to One’s Perceived Experience

Even though the concept of equanimity is not explicitly introduced in Western psychoanalytical theory, it can be found as an important, rather implicit concept within the psychoanalytical approach, where it is believed that (1) it is important for emotions to be seen by the patient with an *even-minded* attitude, without condemning them and (2) the therapist should welcome the patient’s free associations by encountering them with free-floating attention. Psychoanalysis is focused on acknowledgement and acceptance of emotions. For example, Sigmund Freud advised psychoanalysts to instruct patients to “report to us whatever internal perceptions... feelings, thoughts, memories [without] exclud[ing] any of them, whether on the grounds that it is too disagreeable or too indiscreet... or nonsensical” (Freud 1966, p. 287). Freud’s radical proposal was that, without therapy, thoughts or feelings that are unacceptable to the individual are either repressed or suppressed into the unconscious, or projected onto another person (Freud 1966). These defense strategies of “hiding” emotionally challenging experiences have detrimental psychological consequences, since they still exist and can cause psychopathological symptoms. In line with this view, Swiss psychiatrist Carl Jung reportedly declared that

“the foundation of all mental illness is the unwillingness to experience legitimate suffering.” This view is supported by many years of psychology research suggesting that “experiential avoidance,” which can be conceptualized as unhealthy efforts to escape and avoid emotions, thoughts, memories, and other private experiences, contributes to the development and maintenance of many forms of psychopathology (Hayes, Wilson, Gifford, Follette, and Strosahl 1996; Purdon 1999). So while the early psychoanalysts did not promote equanimity practices in the form of formal meditation, the idea that all forms of emotions should be held in awareness with an attitude of acceptance in order to be healthy is the essential foundation of psychoanalytic theory.

While psychoanalysis focuses on the process of bringing unconscious emotions into awareness during the therapeutic encounter, more recent developments in psychotherapy more explicitly teach patients to alter their relationships to their inner experience in general. Buddhist theory has critically inspired “third wave” forms of cognitive and behavioral therapies such as Acceptance and Commitment Therapy (ACT), Dialectical Behavioral Therapy (DBT), and Mindfulness-based Cognitive Therapy (MBCT). In this clinical context, equanimity is still not explicitly mentioned, but is implied in the form of a certain set of attitudes on how to relate to experience, which can be taught in the form of contemplative practices during the therapeutic encounter. These attitudes include nonjudging, non-aversion, non-attachment (or letting-go), and acceptance (Kabat-Zinn 1990; Segal, Williams, and Teasdale 2001). A nonjudging attitude consists in “assuming the stance of an impartial witness to one’s own experience,” which involves suspending judgment and simply “watching whatever comes up,” including observing one’s automatic judgments of “like,” “dislike,” and “don’t-care,” without “judging the judging.” It also involves recognizing one’s thoughts as “just thoughts” without pursuing them any further. Acceptance is the “willingness to see things as they are” (Kabat-Zinn 1990), “not merely tolerance” but rather “the active nonjudgmental embracing of experience in the here and now,” with “undefended ‘exposure’ to thoughts, feelings, and bodily sensations as they are directly experienced” (Hayes, Follette, and Linehan 2004). Here, “exposure” is understood in a similar fashion as in exposure therapy, in that it aims at extinguishing conditioned associations. This exposure must be “undefended,” i.e., in the absence of habitual defense mechanisms such as avoidance or suppression, which are more costly (e.g., Alberts, Schneider, and Martijn 2012) and less effective in the long term (Hayes et al. 2004). The goal is to revert the domination of habitual responses, verbal rules, and evaluations over here-and-now experience, and thereby to facilitate adaptive, flexible responding to environmental contingencies—as opposed to more rigid, rule-governed patterns of responding that are not based in current circumstances (Borkovec 2006; Hayes, Strosahl, and Wilson 1999).

The attitudinal factors listed above are conceptually related to each other (Kabat-Zinn 1990, p. 33), and have been operationalized as a single component of “Attitude” in multicomponent psychological models of mindfulness (Bishop et al. 2004; Shapiro et al. 2006). This component can be summarized as “adopting a particular orientation toward one’s experiences in the present moment, an orientation that is characterized by curiosity, openness, and acceptance” (Bishop et al. 2004). We propose that this attitudinal component corresponds to a shift in perspective toward all experiences or objects, regardless of their affective valence (pleasant, unpleasant, or neutral) or source.

This shift in perspective, in which one watches one’s thought process as if from the outside, like a “fly on the wall” (Ayduk and Kross 2010), facilitates insight towards one’s inner mental landscape and habitual patterns of behavior and has long been acknowledged as an important aspect of psychotherapy in general (Beck et al. 1979). This process has been called psychological self-distancing, decentering, cognitive defusion, re-perceiving, dereification, or metacognitive insight (Bishop et al. 2004; Chambers et al. 2009; Mason and Hargreaves 2001; Shapiro et al. 2006; Teasdale et al. 1995; Teasdale 1999). For example, Safran and Segal (1990) define decentering as the process of observing one’s thoughts and feelings as passing events in the mind, rather than as valid reflections of reality or central aspects of the self. Similarly, Alford and Beck (1997) describe distancing as “the ability to view one’s own thoughts (or beliefs) as constructions of ‘reality’ rather than as reality itself,” and identify this ability as a critical prerequisite for enabling effective therapeutic change. Overall, psychological self-distancing can be said to change one’s relationship to thought rather than attempting to alter the content of thought itself (Hayes et al. 1999).

This perspective shift in relation to one’s experience is by no means limited to psychotherapy or mindfulness practice, but rather is an extension of a fundamental aspect of psychological development and growth across the lifespan (Kegan 1982), also known as “the observing self” (Deikman 1983) or as “meta-awareness” (Chin and Schooler 2010; Schooler et al. 2011). This process allows the individual to uncouple the sensory, directly experienced self from the “narrative” self—the reflective process that maintains continuity of identity across time (Gallagher 2000; Northoff and Bermpohl 2004; Northoff et al. 2006; Williams 2010). In the absence of meta-awareness, the narrative self and other pervasive cognitive processes of “mental simulation” tend to aggravate emotional response (e.g., in the form of prolonged sympathetic arousal), cognitive elaboration, or rumination (Williams 2010). These tendencies are reminiscent of the Buddhist notion of “grasping” or “clinging” mentioned earlier (Analayo 2003; Bodhi 2000) and can become pathological. For example, poor meta-awareness skills increase the risk of recurrence of major depression (Teasdale et al. 2002). Conversely, psychological

self-distancing has been shown to have beneficial effects such as facilitating adaptive self-reflection, decreasing emotional and physiological reactivity, and protecting the individual from negative outcomes associated with rumination over time (Ayduk and Kross 2008, 2010; Kross et al. 2005; Kross and Ayduk 2008).

It has been proposed that mindfulness practice promotes and accelerates the development of this shift in perspective towards an “observing attitude” and yields greater capacity for objectivity about one’s own internal experience and a disidentification from the content of one’s mind (e.g., thoughts, feelings, self-concepts, and memories) (Kerr et al. 2011; Shapiro et al. 2006). This process of disidentification results in greater clarity, perspective, and objectivity (Shapiro et al. 2006). In summary, we propose that this change of perspective—in relation to one’s perceived experience—is a fundamental aspect of equanimity.

Equanimity: An Emotion Regulation Strategy

In concert with the cultivation of an impartial attitude in response to all experience, equanimity involves the skillful process of “maintaining calm and mental equilibrium in the face of provocative stimuli” (Carmody et al. 2009). Such a process fits into traditional conceptualizations of emotion regulation. Emotion regulation (also called self-regulation) refers to any process that influences the onset, offset, magnitude, duration, intensity, or quality of one or more aspects of the emotional response (Gross 1998). Here, we describe equanimity as an emotion regulation strategy that can change both the magnitude and quality of responses.

Different kinds of emotion regulation strategies have been identified (Gross and Thompson 2006; McRae, Ochsner, and Gross 2010; Ochsner and Gross 2005; Parkinson and Totterdell 1999). Based on findings from cognitive neurosciences, Gross and colleagues have suggested a distinction between *behavioral* emotion regulation and *cognitive* emotion regulation (Ochsner and Gross 2005). Behavioral strategies can be antecedent-focused, in which case the appraisal process is controlled through selection or modification of context to avoid the emotion altogether or to modify the emotional impact (e.g., by performing a secondary distracting task). Behavioral strategies can also be response-focused, in which case it is the emotional expression (e.g., smiling, frowning) that is regulated. Cognitive strategies include stimulus–response reversal/extinction, reappraisal (i.e., the reinterpretation of the meaning of a stimulus to change one’s emotional response to it), and attentional control. Of note, these strategies all involve some form of effortful control, unlike equanimity.

While equanimity has some overlap with the emotion regulation strategies described above, it is not identical with them, and in some cases may be opposite (Chambers et al. 2009). For example, in the taxonomy of Ochsner and Gross

(2005), attentional control is regarded as an emotion regulation strategy when attention is guided *away* from the emotion-triggering stimulus. Importantly, while attentional control is also a central aspect of mindfulness practice, in this case the focus of attention is not guided away from the emotion-triggering stimulus, but rather *toward* the stimulus, and also toward the internal emotional experience in response to it—such as the habitual experiences of wanting positive states to last, negative states to end, and neutral states to be less boring (Williams 2010). Mindfulness practice involves bringing sustained attention to the body and resting awareness on sensation during an emotional experience, which leads to a situation of exposure. Equanimity leads to nonreactivity, which then naturally prevents habitual maladaptive response behavior. Engaging the emotional stimulus in this manner regulates the magnitude and duration of the emotional response. Rather than being stuck in the habitual reactions to the external and internal environment, one can experience the transitory nature of all related perceptions, emotions, or cognitions in each moment of experience. Once the purely sensory aspects of the emotional stimulus are recognized, one can choose to disengage from the mental elaborations that usually follow it. In contrast to the response-focused reappraisal strategy, equanimity does not modify the cognitive interpretation of an external stimulus, but rather facilitates the recognition of such thoughts as transient, insubstantial mental events rather than accurate representations of reality (Chambers et al. 2009). The awareness of the transitory nature of one's momentary experience contributes to a dispositional perspective change in relation to one's thoughts. In other words, narrative, self-referential processing (i.e., the narrative of the relevance of the stimulus for oneself) becomes diminished, while the present-moment experience of sensory and mental events becomes enhanced. Thus, continued equanimity practice is predicted to yield the same outcome as stimulus-response reversal methods, where stimuli that formerly provoked reactivity are being responded to with an equanimous state and habitual patterns of maladaptive behavioral response are extinguished.

Equanimity can also indirectly promote cognitive flexibility in response to new and unexpected conditions, such that an optimal response can be chosen based on the current situation, instead of getting “stuck” on habitual thought patterns that may not be the most adaptive response. Cognitive flexibility has been proposed as an indicator of good psychological health (Kashdan and Rottenberg 2010). Both mental balance and cognitive flexibility are important for effective management of stressful emotional events. A balanced state of arousal between hyperexcitability and torpor helps conserve expenditure of mental and physical energy in the face of emotional challenge, while cognitive flexibility broadens the repertoire of possible behavioral responses. Both skills can be enhanced by mindfulness training (Moore and Malinowski 2009).

Equanimity Impacts the Time Course of Emotional Responses

Thoughts and emotions in response to a real or imagined stressor stimulate a coordinated set of bodily responses involving the interaction between the nervous, endocrine, and immune systems (McEwen 1998; Sternberg 2000). Under ideal circumstances, these coordinated responses allow the person to effectively cope with stress, such that when the stressful stimulus has passed, the body gradually returns to its regular state of rest. However, if there is a failure in coping, habituation, or recovery (e.g., blunted or perseverative responses), long-term damage may occur—a phenomenon known as “allostatic load” (Karatsoreos and McEwen 2011; McEwen and Seaman 1999; McEwen 1998, 2008; Sterling and Eyer 1988). Equanimity can aid in the recovery from emotional or physical stress, helping the individual return rapidly back to a state of balance.

As indicated above, the timing of recovery from a stimulus is a crucial component of emotion regulation. Many strong emotions have evolved to function as signaling mechanisms sensitive to the onset and offset of environmental threats towards the individual. These mechanisms have had clear advantages for survival during the course of evolution; however, they are not necessarily adapted to the circumstances of modern life, in which they can lead to chronic stress.

It has been proposed that individuals differ in their emotional style, and that these individual differences can manifest as variations in physiological responses to emotional challenge (Davidson 2003). In Davidson's six-dimensional framework of emotional style, how quickly, or slowly one recovers from stressors corresponds to the “Resilience” dimension (Davidson and Begley 2012). Davidson (2003) proposed that specific parameters of physiological response to emotional challenge can be objectively measured, including the threshold to respond, the magnitude of the response, the rise time to the peak of the response, the recovery function of the response, and the duration of the response (see Fig. 1). The last three parameters correspond to different aspects of affective chronometry or time course for emotional responding (Davidson 2003). We propose that these parameters could be used to measure certain aspects of equanimity. For example, one could hypothesize that increased equanimity manifests as a reduction in magnitude and/or duration of certain emotions (especially those considered maladaptive), while not affecting other ones, or even amplifying some of them: in the context of specific Buddhist meditation practices aimed at cultivating compassion for others' suffering, the magnitude of emotional response was found to increase in expert meditators, who presumably have high equanimity (Lutz et al. 2008).

We propose that a primary “signature” of equanimity is in the temporal domain, in the form of a more rapid disengagement from initial emotional response and faster return to baseline. For example, in response to a loud noise, there




Fig. 1 Generic representation of a typical physiological response to an emotional stimulus. Different parameters of affective style and chronometry are depicted. The magnitude of response corresponds to the height of the response at its peak. Duration is the time from the onset of response to

the return to baseline. In this example, an equanimous response involves a rapid recovery function (*solid line*), in contrast to one that is either low in magnitude (“blunted” *dotted line*) or prolonged (“perseverative” *dotted line*)

may be a similar initial startle response, but with equanimity it is less likely to be followed by anxiety and the person will quickly return to their regular state. Furthermore, in our opinion, equanimity does not necessarily entail a complete lack of physiological response to emotional challenges. A lack of emotional response in circumstances when it is normally expected is usually considered a mark of psychopathology. As discussed above, mere indifference is very distinct from equanimity.

Measuring Equanimity

Assessing equanimity, both subjectively and objectively, presents several challenges. For instance, as discussed above, equanimity may be difficult to distinguish from apathy, as both can lead to diminished emotional responses to a given stimulus or context. However, in the case of equanimity, one still experiences care and interest in the situation. Therefore, a scientific measure of equanimity will ideally be able to distinguish it from apathy. Below, we review previous studies that have used a variety of psychological, physiological, and neuroimaging methods to assess equanimity or emotional responding, either in experienced meditation practitioners or in longitudinal studies with novices, and suggest ways that these measures could be further developed towards a more complete measure of equanimity.

Self-report Measures of Equanimity

A number of researchers have proposed self-report scales that target equanimity in the sense of resilience or effective management of stressful life events (Büssing et al. 2006; Kraus and Sears 2009; Lundman et al. 2007; Mack et al. 2008; Van Tongeren and Green 2010); however, most scales lack a theoretical framework and reveal the lack of a common agreement for the construct of equanimity.

Kraus and Sears (2009) developed a questionnaire to measure the four “immeasurable qualities” at the heart of Buddhist teachings: loving-kindness, compassion, appreciative (or sympathetic) joy, and equanimity. However, equanimity was only represented by one item, namely “acceptance.”

Mack et al. (2008) developed the Peace, Equanimity, and Acceptance in Cancer Experience (PEACE) scale. Equanimity was not explicitly defined but items on this questionnaire address “acceptance of diagnosis,” “sense of inner peace and harmony,” “having made peace with it,” and “sense of inner calm and tranquility.”

Büssing et al. (2006) developed a questionnaire to measure distinct expressions of spirituality. One of the seven factors is equanimity, which is comprised of three items (“trying to practice equanimity,” “trying to achieve a calm spirit,” and “meditate”). These items are measuring the subject’s attempts or efforts towards equanimity, rather than measuring equanimity directly.

In an assessment of the psychometric properties of a Resilience Scale, Lundman et al. (2007) conceptualized equanimity as one of the factors of resilience. They define equanimity as “a balanced perspective of one’s life.” Sample items of this scale include: “I take things 1 day at a time,” “I do not dwell on things that I can’t do anything about,” “I usually take things in stride,” “I seldom wonder what the point of it all is,” “It’s okay if there are people who don’t like me,” and “When I’m in difficult situations, I can usually find my way out of it.” Not only do these questionnaire items conflate multiple constructs (such as self-worth and rumination), but the conceptualization of equanimity overall does not differentiate equanimity from apathy.

Van Tongeren and Green (2010) propose that psychological equanimity is related to the integrity of the self or the integrity of one’s own world views. They suggest that in order to maintain psychological equanimity, sources of meaning have to be preserved. Whenever an individual experiences disruption to one domain of self-meaning, alternative sources

of meaning are being bolstered to regain equanimity. This understanding of the term suggests an evaluative process equating equanimity with a state of alignment or balance between one's world view and one's personal experience. This definition is at odds with the one that we propose here based on the Buddhist tradition.

Other scales have recently been developed that assess constructs closely related to equanimity. In the Nonattachment Scale (NAS), informed by classic and contemporary Buddhist scholarship, nonattachment is described as psychological flexibility, nonreactivity, mindedness, more quickly recovering from upsets, allowing, releasing, supporting others' capacity to choose, and a sense of ease (Sahdra et al. 2010). Phenomenologically, nonattachment is described as having the subjective quality of not being stuck or fixated on ideas, images, or sensory objects and not feeling an internal pressure to acquire, hold, avoid, or change. This description corresponds well to the Buddhist definition of equanimity we have described here. Example items include: "I can accept the flow of events in my life without hanging onto them or pushing them away," "I can let go of regrets and feelings of dissatisfaction about the past," and "I can remain open to what life offers me regardless of whether it seems desirable or undesirable at a particular time." Sahdra et al. (2010) address the distinction between nonattachment and indifference and demonstrate that the scale is inversely related to avoidant attachment, dissociation, alexithymia, and impersonal motivational orientation. While the scale includes an array of related constructs (e.g., the willingness to view difficulties as challenges and the capacity to take joy in others' achievements), we feel that many of its items would be well-suited to assess the subjective aspects of equanimity.

Another construct that is related to equanimity is the concept of decentering. Earlier, we elaborated on how this change in relation to one's experience is essential for the development of equanimity (see "[Equanimity: A Relation to One's Perceived Experience](#)"). The Experience Questionnaire developed by Teasdale, Segal, and Williams assesses decentering, along with rumination. Studies have demonstrated good psychometric properties and construct validity of the scale and showed that the relevant items load on a single factor of decentering (Fresco et al. 2007). The construct is negatively related to experiential avoidance and positively related to emotion regulation. Sample items include "I can observe unpleasant feelings without being drawn into them" and "I can separate myself from my thoughts and feelings." Future research could employ this scale to test the assumption that decentering is an important process for developing equanimity, and could be used to further reveal the relationship between the two constructs.

Finally, it should be noted that any self-report measure of equanimity is likely to face the same problems that the existing mindfulness questionnaires are facing, namely: (1) experts may disagree about what the construct is; (2)

questionnaire designers may lack a deep theoretical and experiential understanding of the construct; (3) items may be differently interpreted, depending on the development and meditation practice of respondents; (4) there may be discrepancies between self-report and reality; (5) biases may occur in meditation practitioners due to their large investment of time and effort in the practice; and (6) the validation of instruments may be compromised by choosing inappropriate populations (Grossman 2008). While a number of these issues can be addressed, some are inherent to all self-report measures.

Physiological Measures Indicative of Equanimity

In addition to the inherent limitations of self-report measures, study participants may not recognize their own equanimity if some emotions are still present, albeit transient and short-lived. Some of these emotions may even have a high intensity, especially in individuals who had a tendency for blunted emotions before meditation training. Therefore, we propose that objective measures should be pursued to provide a more meaningful and accurate metric of equanimity. Physiological measures that may be relevant to measure equanimity objectively include autonomic, endocrine, and inflammatory markers.

Indices of autonomic function have been used for decades in the field of psychophysiology to assess how emotions manifest in the body (Ledoux 1996). Popular indices include skin conductance, heart rate, respiratory rate, and heart rate variability (HRV). For example, skin conductance varies with sympathetic nervous system activation and can be used as an indication of psychological or physiological arousal. It is well known that the "fight-or-flight" response elicits increases in heart rate and breathing rate, indicative of higher sympathetic activation (Cannon 1915). Conversely, the "relaxation response" is characterized by a decrease in heart rate and breathing rate and an increase in HRV, consistent with increased parasympathetic activation (Benson 1975). Many studies to date have investigated how these autonomic indices relate to emotion regulation (e.g., Chapman et al. 2010; Demaree et al. 2004; Dikecligil and Mujica-Parodi 2010; Geisler et al. 2010; Gyurak and Ayduk 2008; Ohira et al. 2006; Porges 2007, 2011; Pu et al. 2010; Ruiz-Padial et al. 2011; Smith et al. 2011; Thayer and Lane 2000; Thayer et al. 2012; Vohs and Baumeister 2010; Volokhov and Demaree 2010; Campbell-Sills et al. 2006; Driscoll et al. 2009; Najström and Jansson 2007; Robinson and Demaree 2009; Sheppes et al. 2009). We propose that these indices may also be used to assess equanimity from a physiological perspective. For example, one component of HRV (namely, respiratory sinus arrhythmia, which measures vagal cardiovascular reactivity) was used as a measure of emotional equanimity in a study on emotional and motivational orientation towards academic performance in college students (Gramzow et al. 2008). Emotional

equanimity in this context was defined as a “calm and composed state” and as “keeping a cool head when thinking about academic performance (i.e., a challenge).” Another potentially relevant index is heart rate deceleration, a physiological response that has been associated with orienting toward and engaging with others (Dunn et al. 2009; Eisenberg et al. 1989; Goetz et al. 2010; Stellar et al. 2012).

Other physiological markers relevant to emotion include stress hormones and immune markers such as cortisol and inflammatory cytokines, which affect the brain and are influenced by it in a bidirectional fashion (Sternberg 2000). These markers are sometimes measured during an emotional or stress challenge such as the Trier Social Stress Test (TSST), in which the subject is asked to deliver a speech and perform a challenging mental arithmetic task in front of an audience (Kirschbaum et al. 1993). The TSST induces increases in cortisol and inflammatory cytokines over the course of several hours following the stressor (Steptoe et al. 2007; Yamakawa et al. 2009). However, responses vary across individuals, and it has been proposed that this variability may correspond to differences in emotion regulation skills across individuals (Pace et al. 2006; von Känel et al. 2006; Wirtz et al. 2007). In the context of a stress test, equanimity could be operationalized as a lower overall level of stress hormones and/or a shorter time course for the return to baseline, presumably mirroring subjects’ inner process of self-distancing and decentering. For example, Pace et al. (2009) found a post hoc association between compassion meditation practice time and decreased levels of interleukin-6, a proinflammatory cytokine, although there was no difference between the meditation group and the control group overall. In other work, Hoge and colleagues found lower levels of adrenocorticotrophin hormone (the hormone that stimulates cortisol release) during the TSST in a Mindfulness-Based Stress Reduction (MBSR) group, compared to an active control group (Hoge et al. 2012).

Another measure of emotion response that could be applied to the study of equanimity is the activation of the facial muscles involved in facial expressions (Ekman and Rosenberg 2005). In one case study, an experienced Buddhist monk showed a smaller startle response to a loud noise (as measured by electrical activity of the muscles controlling eye blinks) during meditation than during distraction. In addition, the startle response was smaller in a meditative state of equanimity called “Open Awareness” (*rig pa cog gzhas* in Tibetan) than in a meditative state of focused concentration on an external object (Levenson et al. 2012). This case study suggests that external physiological measures can be indicative of different levels of equanimity.

The above two sections describe complementary methods that could be used to measure equanimity. While physiological measurements may give more objective assessments, supplementary psychological self-reports could be used in combination with laboratory measures to help distinguish

equanimity from apathy or other forms of top-down emotional regulation (e.g., reappraisal, suppression, or distraction), which may also manifest as a lower-amplitude physiological response to a stimulus. After meditation training, subjects tend to provide more detailed phenomenological reports of their moment-by-moment experience, which can be of great value for the scientific investigation of mental processes (Varela et al. 1992).

Putative Neural Mechanisms of Equanimity

A number of recent neuroimaging studies have revealed possible neural mechanisms underlying equanimity, using various forms of emotional challenges, including pain. While these studies were not explicitly designed to investigate equanimity per se, we believe that they offer important insights into how equanimity might manifest in the brain in various experimental contexts.

It was previously found that compared to novices, long-term meditators had a significant reduction of self-reported unpleasantness—but not intensity—of painful stimuli while practicing open-presence meditation (Perlman et al. 2010). In a recent study of experienced Vipassana meditators, these findings were replicated and extended with a neuroimaging investigation of the underlying neural mechanisms (Gard et al. 2012). The reduction in pain unpleasantness while in a meditative state of mindfulness was accompanied by increased brain activation in sensory areas (the posterior insula/secondary somatosensory cortex) and decreased activation in an area that mediates executive control (the lateral prefrontal cortex). This suggests that this meditative state corresponds to a decrease in cognitive control of the pain, rather than a decrease in the pain sensation per se. Another study investigated pain processing in experienced Zen meditators who were explicitly instructed *not* to enter a meditative state (Grant et al. 2011). In response to pain (heat) stimulation, the experienced meditators exhibited lower pain sensitivity together with greater activation in primary pain sensory processing regions (insula, thalamus, and anterior cingulate cortex) and less activation in executive, evaluative, and emotion-related areas (prefrontal cortex, amygdala, and hippocampus) compared to controls. Importantly, the lower pain sensitivity in meditators was strongly predicted by decoupling between activity in the cognitive–evaluative prefrontal areas and sensory pain areas (Grant et al. 2011). This decoupling may correspond to the neural substrate of the meditators’ capacity to view painful stimuli more neutrally, which is one aspect of equanimity.

Other neuroimaging studies have investigated the neural correlates of emotion regulation in the context of meditation. As reviewed below, these studies point to an important, if complex, role of the amygdala—a brain region centrally involved in the perception and appraisal of emotional (negative as well as positive) stimuli and in orienting attention towards

emotionally significant stimuli (Vuilleumier 2005; Whalen and Phelps 2009; Whalen 1998). Previous studies indicate that emotional regulation strategies such as cognitive distancing and reappraisal promote decreased amygdala activation (Beauregard et al. 2001; Lévesque et al. 2003; Ochsner et al. 2002, 2004), and conversely, the instruction to “maintain” the emotional state elicited by negative-valence stimuli causes an increase in amygdala activation (Schaefer et al. 2002).

In one of several recent neuroimaging studies of meditation, experienced Zen meditators perceived emotional pictures as less intense while being in a state of mindfulness than during rest, but that this happened without increased activity in modulatory prefrontal brain areas and without downregulation of the amygdala, suggesting that it was not just a standard case of top-down control over low-level affective cerebral systems from higher-level cortical brain regions (Taylor et al. 2011). In contrast, in the same study, novice meditators (20 min/day for 7 days) instructed to be in a state of mindfulness showed increased activation in regulatory prefrontal brain areas and decreased activation in the amygdala when successfully downregulating emotional intensity. The difference between experienced and novice meditators indicates that emotion regulation strategies may shift dramatically with extensive meditation experience, following the development of greater equanimity that no longer requires active downregulation of emotions.

Another recent functional magnetic resonance imaging (fMRI) study suggests that, after 8 weeks of *shamatha* (focused attention) meditation training, novice meditators exhibited a longitudinal decrease in amygdala activation in response to emotion-evoking stimuli, compared to before training (Desbordes et al. 2012). Importantly, throughout the experiment subjects remained in an ordinary, nonmeditative state. This finding suggests that 8 weeks of meditation training may affect emotional processing in everyday life, and not just during meditation. In another recent fMRI study, generalized

anxiety disorder patients exhibited a reduction in amygdala activation in response to neutral facial expressions following participation in an MBSR course (Hölzel et al. 2013). In addition, these patients also showed altered functional connectivity between the amygdala and regions of the prefrontal cortex which are known to be involved in the monitoring of arousal. While studies on emotion regulation typically show a negative correlation between prefrontal regions and the amygdala when subjects downregulate emotional responses, this study found a positive correlation between amygdala and prefrontal activation following mindfulness training, suggesting that the typical downregulation of reactivity through the prefrontal cortex may have been replaced with a mere monitoring of arousal (amygdala activation) through the prefrontal cortex. These findings appear in line with our conceptualization of equanimity as the observance rather than suppression of emotional responses.

In line with the above findings, patients with social anxiety disorder who had completed an 8-week MBSR course showed decreased negative emotional experience, decreased amygdala activation, and decreased activation in other brain regions involved in attention deployment, when contrasting a condition of mindful breath awareness with a condition of reacting to phrases expressing negative self-beliefs (Goldin and Gross 2010). Interestingly, this study also showed a longitudinal difference in the timing of amygdala activation during the initial reaction to negative self-belief phrases (a form of emotional challenge), when comparing before and after MBSR. Before MBSR training, the amygdala response to negative self-belief phrases slowly rose throughout the 12-s stimulus presentation (see Fig. 2, “React” phase). After MBSR training, the amygdala response initially rose quickly, but then reverted after 4.5 s and decreased significantly, several seconds before the subjects were instructed to enter a state of mindful breath awareness (see Fig. 2). This finding is in agreement with our proposition of a faster return to baseline in response to




Fig. 2 Right dorsal amygdala blood oxygenation level-dependent (BOLD) contrast signal time series during reacting to negative self-beliefs and breath-focused attention in social phobics (SP) at both pre- and post-

MBSR. $*p < 0.05$. *Rate* negative emotion rating, *React* reacting to the negative self-belief, *Breath-Focus* instruction to focus attention on breath sensation. Reproduced from Goldin and Gross (2010)

emotional stressors that would correspond to greater equanimity. Of note, the time course of amygdala activity in response to negative emotion stimuli has been linked to trait measures of affective style: Schuyler et al. (2012) found that a slower amygdala recovery was associated with greater trait neuroticism, which is characterized by a decreased ability to regulate emotion and greater perseveration following negative events (Kokkonen and Pulkkinen 2001; Suls and Martin 2005), i.e., low equanimity as defined here.

A few other studies have investigated the temporal aspect of emotional reactivity and engagement with emotion. One study employing an emotional interference task showed that participants of a mindfulness intervention disengaged more rapidly from the unpleasant interfering stimulus than participants of a relaxation intervention or waitlist controls (Ortner et al. 2007). In a sample of female fibromyalgia patients, a preliminary study of mindfulness meditation on attentional bias revealed decreased avoidance at early stages of attention towards pain-related cues and more rapid disengagement from threat at later stages of cognitive processing in those patients exposed to 8 weeks of mindfulness training (Vago and Nakamura 2011). In another study, Zen practitioners had a reduced duration of neural response associated with conceptual processing in regions of the default network as compared to matched controls during a lexical decision task, suggesting reduced elaboration or reactivity to stimuli (Pagnoni et al. 2008). In addition, the time course of activation in the brain's attention network was more strongly correlated with the time course of the task itself in meditators than in controls (Guo and Pagnoni 2008). These studies indicate that subjects trained in meditation show higher equanimity as defined by a faster disengagement from a stimulus once it has become irrelevant.

Most studies discussed above indicate that experienced meditators, who purportedly have high levels of equanimity, show decreased activation in evaluative and regulatory brain regions but increased activation in sensory brain regions in response to emotion-evoking stimuli. The above studies suggest that while experienced meditators exhibit neural activation patterns indicative of equanimity, beginner meditators tend to show patterns reflecting increased cognitive control and regulation. This is consistent with our hypothesis above that equanimity may take a longer time to develop than purely attentional skills. In this regard, we should note that it would be particularly beneficial for future studies to include expert meditators as full-fledged collaborators (rather than simple experimental subjects) as their personal experience with meditation practice may provide invaluable information towards study design and hypotheses—especially if they are also trained in scientific methods (Desbordes and Negi 2013).

Beyond the emotional response to specific stimuli, recent neuroimaging studies have begun to look at spontaneously occurring brain states in expert meditators, who may have overall greater equanimity than nonmeditators. The

investigation of spontaneous brain activity is a particularly promising approach for studying equanimous states because it may reveal underlying changes in connectivity between brain regions and synchrony within neuronal assemblies that are thought to shape our moment-by-moment responses to stimuli. For example, Brewer et al. (2011) found lower activation in the brain's default mode network, which is associated with mind wandering and self-referential processing, in experienced meditators compared to meditation-naïve controls. In addition, the same group found that, in experienced meditators, deactivation in a core region of the default mode network (the posterior cingulate cortex) was related on a moment-by-moment basis to subjective reports of “not efforting,” “contentment,” and other descriptions consistent with the experience of an equanimous state (Garrison et al. 2013a, b). Other recent studies also found differences in functional connectivity in the default mode network between meditators and controls (Jang et al. 2011; Kilpatrick et al. 2011). Overall, these findings are consistent with the hypothesis that the cultivation of specific meditative states, which are relatively short-term, can result in enduring changes in mental function, i.e., in the long-term development of certain traits such as equanimity (Slagter et al. 2011).

Conclusions and Future Directions

In summary, we introduced equanimity as a fundamental concept in meditation practice that we propose captures an essential psychological element in the improvement of well-being, and therefore should be a focus in future research studies of meditation and other contemplative practices.

Dissociating equanimity from mindfulness is a critical step in dismantling the neurocognitive, psychological, physiological, and behavioral mechanisms that enable the transformation of maladaptive processes into more adaptive ones. Here, we describe equanimity as a state and dispositional tendency that can be developed over time through specific contemplative practices. Equanimity transforms our sensory–perceptual and cognitive–emotional systems to widen our perspective on experience, more readily engage incoming sensory information, and more efficiently disengage cognitive–evaluative and emotional–reactive behaviors when appropriate. Generally, this transformation is more gradual than the development of enhanced meta-awareness. Identifying mindfulness and equanimity as separate mental factors also facilitates a more cohesive and accurate nomenclature. Terms such as “distancing,” “accepting,” “non-judging,” and “non-attachment” describe certain aspects of equanimity, but are incomplete and often misleading, causing confusion and misunderstanding in the psychological literature.

Understanding the distinction between mindfulness and equanimity will also facilitate the development of more precise quantitative tools to assess each. Given the significant

difficulties with previous self-report psychometrics in meditation research, we also propose that equanimity should be measured objectively in the laboratory instead of constricting this dynamic process into a solely self-report questionnaire. While a self-report component will be useful for developing and validating the interpretation of objective metrics such as brain activity or physiological markers, most individuals will not be able to accurately assess their own levels of equanimity. In particular, in some instances the emotional response to a given stimulus may become larger, though more transient, with the development of equanimity, which could be confusing and counter-intuitive for many novice practitioners. It should also be noted that although we have described how equanimity can be developed through the practice of meditation techniques, this does not exclude the development or expression of equanimity within other contexts, such as traditional psychotherapeutic techniques or mind-body practices (e.g. yoga, tai-chi, etc.). We hope this article will stimulate research into developing new laboratory measures of equanimity that more comprehensively capture this important, multidimensional construct.

Perhaps this is a small contribution to a larger shift in orientation around the whole issue of well-being and human flourishing, which so often seems to aim at maximizing pleasure and sustaining happiness while avoiding or eliminating pain and ameliorating the effects of emotional discomfort. It may be that overall well-being is better served by learning to cultivate equanimity, both as an effective state for responding skillfully to whatever is arising in the present moment, and as a healthy trait that can be strengthened over time and integrated into one's character.

Acknowledgments This work was supported in part by the National Institutes of Health (grant number K23 AT004432 to Elizabeth Hoge) and the 7th European Community Framework Programme (Marie Curie International Outgoing Fellowship to Britta K. Hölzel). The authors would like to thank Andrea Grabovac for helpful comments on an earlier version of this manuscript. All authors contributed equally to this work and are listed alphabetically.

References

- Alberts, H. J. E. M., Schneider, F., & Martijn, C. (2012). Dealing efficiently with emotions: acceptance-based coping with negative emotions requires fewer resources than suppression. *Cognition & Emotion*, *26*(5), 863–870.
- Alford, B. A., & Beck, A. T. (1997). *The integrative power of cognitive therapy*. New York: The Guilford Press.
- Analayo, (2003). *Satipatthana: the direct path to realization*. Cambridge: Windhorse.
- Austin, J. H. (2009). *Selfless insight: Zen and the meditative transformations of consciousness*. Cambridge: MIT Press.
- Ayduk, Ö., & Kross, E. (2008). Enhancing the pace of recovery: self-distanced analysis of negative experiences reduces blood pressure reactivity. *Psychological Science*, *19*(3), 2292–2231.
- Ayduk, Ö., & Kross, E. (2010). From a distance: implications of spontaneous self-distancing for adaptive self-reflection. *Journal of Personality and Social Psychology*, *98*(5), 809–829.
- Baer, R. A., Smith, G. T., Lykins, E. L., Button, D., Krietemeyer, J., Sauer, S., et al. (2008). Construct validity of the five facet mindfulness questionnaire in meditating and nonmeditating samples. *Assessment*, *15*(3), 329–342.
- Beauregard, M., Lévesque, J., & Bourgouin, P. (2001). Neural correlates of conscious self-regulation of emotion. *Journal of Neuroscience*, *21*(18), RC165.
- Beck, A. T., Rush, A. J., Shaw, B. F., & Emery, G. (1979). *Cognitive therapy of depression*. New York: The Guilford Press.
- Benson, H. (1975). *The relaxation response*. New York: Morrow.
- Bishop, S. R., Lau, M., Shapiro, S. L., Carlson, L., Anderson, N. D., Carmody, J., et al. (2004). Mindfulness: a proposed operational definition. *Clinical Psychology: Science and Practice*, *11*(3), 230–241.
- Bodhi, B. (2000). *A comprehensive manual of Abhidhamma: the philosophical psychology of Buddhism*. Onalaska: Buddhist Publication Society Pariyatti Editions.
- Bodhi, B. (2005). *In the Buddha's words: an anthology of discourses from the Pali Canon*. Somerville: Wisdom Publications.
- Bodhi, B. (2011). What does mindfulness really mean? A canonical perspective. *Contemporary Buddhism*, *12*(1), 19–39.
- Borkovec, T. D. (2006). Life in the future versus life in the present. *Clinical Psychology: Science and Practice*, *9*(1), 76–80.
- Brewer, J. A., Worhunsky, P. D., Gray, J. R., Tang, Y.-Y., Weber, J., & Kober, H. (2011). Meditation experience is associated with differences in default mode network activity and connectivity. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(50), 20254–20259.
- Buddhaghosa, B. (2003). *The path of purification: Visuddhimagga*. Onalaska: Buddhist Publication Society Pariyatti Editions.
- Büssing, A., Ostermann, T., & Matthiessen, P. F. (2006). Distinct expressions of vital spirituality: the ASP questionnaire as an explorative research tool. *Journal of Religion and Health*, *46*(2), 267–286.
- Campbell-Sills, L., Barlow, D. H., Brown, T. A., & Hofmann, S. G. (2006). Effects of suppression and acceptance on emotional responses of individuals with anxiety and mood disorders. *Behaviour Research and Therapy*, *44*(9), 1251–1263.
- Cannon, W. B. (1915). *Bodily changes in pain, hunger, fear and rage: an account of recent researches into the function of emotional excitement*. New York: Appleton.
- Carmody, J., Baer, R. A., Lykins, E. L., & Olendzki, N. (2009). An empirical study of the mechanisms of mindfulness in a mindfulness-based stress reduction program. *Journal of Clinical Psychology*, *65*(6), 613–626.
- Chambers, R., Gullone, E., & Allen, N. B. (2009). Mindful emotion regulation: an integrative review. *Clinical Psychology Review*, *29*(6), 560–572.
- Chapman, H. A., Woltering, S., Lamm, C., & Lewis, M. D. (2010). Hearts and minds: coordination of neurocognitive and cardiovascular regulation in children and adolescents. *Biological Psychology*, *84*(2), 296–303.
- Chin, J., & Schooler, J. W. (2010). Meta-awareness. In W. P. Banks (Ed.), *Encyclopedia of consciousness* (pp. 33–41). San Diego: Elsevier.
- Dalai Lama. (2001a). *Stages of meditation*. Ithaca: Snow Lion.
- Dalai Lama. (2001b). *Dzogchen: the heart essence of the great perfection*. Ithaca: Snow Lion.
- Dalai Lama. (2001c). *An open heart: practicing compassion in everyday life*. New York: Little, Brown and Company.
- Davidson, R. J. (2003). Darwin and the neural bases of emotion and affective style. *Annals of the New York Academy of Sciences*, *1000*(1), 316–336.
- Davidson, R. J., & Begley, S. (2012). *The emotional life of your brain: how its unique patterns affect the way you think, feel, and live—and how you can change them*. New York: Hudson Street Press.
- Deikman, A. J. (1983). *The observing self: mysticism and psychotherapy*. Boston: Beacon.

- Demaree, H. A., Robinson, J. L., Everhart, D. E., & Schmeichel, B. J. (2004). Resting RSA is associated with natural and self-regulated responses to negative emotional stimuli. *Brain and Cognition*, *56*(1), 14–23.
- Desbordes, G., & Negi, L. T. (2013). A new era for mind studies: training investigators in both scientific and contemplative methods of inquiry. *Frontiers in Human Neuroscience*, *7*, 741.
- Desbordes, G., Negi, L. T., Pace, T. W. W., Wallace, B. A., Raison, C. L., & Schwartz, E. L. (2012). Effects of mindful-attention and compassion meditation training on amygdala response to emotional stimuli in an ordinary, non-meditative state. *Frontiers in Human Neuroscience*, *6*, 292.
- Dikeçligil, G. N., & Mujica-Parodi, L. R. (2010). Ambulatory and challenge-associated heart rate variability measures predict cardiac responses to real-world acute emotional stress. *Biological Psychiatry*, *67*(12), 1185–1190.
- Dreyfus, G. (2011). Is mindfulness present-centred and non-judgmental? A discussion of the cognitive dimensions of mindfulness. *Contemporary Buddhism*, *12*(1), 41–54.
- Driscoll, D., Tranel, D., & Anderson, S. W. (2009). The effects of voluntary regulation of positive and negative emotion on psychophysiological responsiveness. *International Journal of Psychophysiology*, *72*(1), 61–66.
- Dunn, B. D., Billotti, D., Murphy, V., & Dalgleish, T. (2009). The consequences of effortful emotion regulation when processing distressing material: a comparison of suppression and acceptance. *Behaviour Research and Therapy*, *47*(9), 761–773.
- Dunne, J. D. (2011). Toward an understanding of non-dual mindfulness. *Contemporary Buddhism*, *12*(1), 71–88.
- Eisenberg, N., Fabes, R. A., Miller, P. A., Fultz, J., Shell, R., Mathy, R. M., et al. (1989). Relation of sympathy and personal distress to prosocial behavior: a multimethod study. *Journal of Personality and Social Psychology*, *57*(1), 55–66.
- Ekman, P., & Rosenberg, E. L. (Eds.). (2005). *What the face reveals: basic and applied studies of spontaneous expression using the Facial Action Coding System (FACS)*. New York: Oxford University Press.
- Fresco, D. M., Moore, M. T., van Dulmen, M. H. M., Segal, Z. V., Ma, S. H., Teasdale, J. D., et al. (2007). Initial psychometric properties of the experiences questionnaire: validation of a self-report measure of decentering. *Behavior Therapy*, *38*(3), 234–246.
- Freud, S. (1966). *The complete introductory lectures on psychoanalysis*. New York: Norton.
- Gallagher, I. (2000). Philosophical conceptions of the self: implications for cognitive science. *Trends in Cognitive Sciences*, *4*(1), 14–21.
- Gard, T., Hölzel, B. K., Sack, A. T., Hempel, H., Lazar, S. W., Vaitl, D., et al. (2012). Pain attenuation through mindfulness is associated with decreased cognitive control and increased sensory processing in the brain. *Cerebral Cortex*, *22*(11), 2692–2702.
- Garrison, K. A., Santoyo, J. F., Davis, J. H., Thornhill, T. A., Kerr, C. E., & Brewer, J. A. (2013a). Effortless awareness: using real time neurofeedback to investigate correlates of posterior cingulate cortex activity in meditators' self-report. *Frontiers in Human Neuroscience*, *7*, 440.
- Garrison, K. A., Scheinost, D., Worhunsky, P. D., Elwafi, H. M., Thornhill, T. A., Thompson, E., et al. (2013b). Real-time fMRI links subjective experience with brain activity during focused attention. *NeuroImage*, *81*, 110–118.
- Geisler, F. C. M., Vennewald, N., Kubiak, T., & Weber, H. (2010). The impact of heart rate variability on subjective well-being is mediated by emotion regulation. *Personality and Individual Differences*, *49*(7), 723–728.
- Goetz, J. L., Keltner, D., & Simon-Thomas, E. (2010). Compassion: an evolutionary analysis and empirical review. *Psychological Bulletin*, *136*(3), 351–374.
- Goldin, P. R., & Gross, J. J. (2010). Effects of mindfulness-based stress reduction (MBSR) on emotion regulation in social anxiety disorder. *Emotion*, *10*(1), 83–91.
- Grabovac, A. D., Lau, M. A., & Willett, B. R. (2011). Mechanisms of mindfulness: a Buddhist psychological model. *Mindfulness*, *2*(3), 154–166.
- Gramzow, R. H., Willard, G., & Mendes, W. B. (2008). Big tales and cool heads: academic exaggeration is related to cardiac vagal reactivity. *Emotion*, *8*(1), 138–144.
- Grant, J. A., Courtemanche, J., & Rainville, P. (2011). A non-elaborative mental stance and decoupling of executive and pain-related cortices predicts low pain sensitivity in Zen meditators. *Pain*, *152*(1), 150–156.
- Gross, J. J. (1998). The emerging field of emotion regulation: an integrative review. *Review of General Psychology*, *2*(3), 271–299.
- Gross, J. J., & Thompson, R. A. (2006). Emotion regulation: conceptual foundations. In J. J. Gross (Ed.), *Handbook of emotion regulation*. New York, NY: The Guilford Press.
- Grossman, P. (2008). On measuring mindfulness in psychosomatic and psychological research. *Journal of Psychosomatic Research*, *64*(4), 405–408.
- Gunaratana, B. H. (2002). *Mindfulness in plain English*. Boston: Wisdom Publications.
- Guo, Y., & Pagnoni, G. (2008). A unified framework for group independent component analysis for multi-subject fMRI data. *NeuroImage*, *42*(3), 1078–1093.
- Gyurak, A., & Ayduk, Ö. (2008). Resting respiratory sinus arrhythmia buffers against rejection sensitivity via emotion control. *Emotion*, *8*(4), 458–467.
- Hayes, S. C., Wilson, K. G., Gifford, E. V., Follette, V. M., & Strosahl, K. (1996). Experimental avoidance and behavioral disorders: a functional dimensional approach to diagnosis and treatment. *Journal of Consulting and Clinical Psychology*, *64*(6), 1152–1168.
- Hayes, S. C., Strosahl, K. D., & Wilson, K. G. (1999). *Acceptance and commitment therapy: an experiential approach to behavior change*. New York: The Guilford Press.
- Hayes, S. C., Follette, V. M., & Linehan, M. M. (Eds.). (2004). *Mindfulness and acceptance: expanding the cognitive-behavioral tradition*. New York: The Guilford Press.
- Hoge, E. A., Bui, T. H. E., Metcalf, C. E., Pollack, M. H., Simon, N. M. (2012). Mindfulness training improves resilience: Reductions in adrenocorticotropic hormone (ACTH) response to laboratory stress. *American College of Neuropsychopharmacology 51st Annual meeting*, Miami, FL, December 5, 2012.
- Hölzel, B. K., Hoge, E. A., Greve, D. N., Gard, T., Creswell, J. D., Brown, K. W., et al. (2013). Neural mechanisms of symptom improvements in generalized anxiety disorder following mindfulness training. *NeuroImage: Clinical*, *2*, 448–458.
- Jang, J. H., Jung, W. H., Kang, D.-H., Byun, M. S., Kwon, S. J., Choi, C.-H., et al. (2011). Increased default mode network connectivity associated with meditation. *Neuroscience Letters*, *487*(3), 358–362.
- Josipovic, Z. (2010). Duality and nonduality in meditation research. *Consciousness and Cognition*, *19*(4), 1119–1121.
- Kabat-Zinn, J. (1990). *Full catastrophe living: using the wisdom of your body and mind to face stress, pain, and illness*. New York: Delacorte Press.
- Kang, C., & Whittingham, K. (2010). Mindfulness: a dialogue between Buddhism and clinical psychology. *Mindfulness*, *1*(3), 161–173.
- Karatsoreos, I. N., & McEwen, B. S. (2011). Psychobiological allostasis: resistance, resilience and vulnerability. *Trends in Cognitive Sciences*, *15*(12), 576–584.
- Kashdan, T. B., & Rottenberg, J. (2010). Psychological flexibility as a fundamental aspect of health. *Clinical Psychology Review*, *30*(7), 865–878.
- Kegan, R. (1982). *The evolving self: problem and process in human development*. Cambridge: Harvard University Press.
- Kerr, C. E., Josyula, K., & Littenberg, R. (2011). Developing an observing attitude: an analysis of meditation diaries in an

- MBSR clinical trial. *Clinical Psychology & Psychotherapy*, 18(1), 80–93.
- Kilpatrick, L. A., Suyenobu, B. Y., Smith, S. R., Bueller, J. A., Goodman, T., Creswell, J. D., et al. (2011). Impact of mindfulness-based stress reduction training on intrinsic brain connectivity. *NeuroImage*, 56(1), 290–298.
- Kirschbaum, C., Pirke, K. M., & Hellhammer, D. H. (1993). The “Trier Social Stress Test”—a tool for investigating psychobiological stress responses in a laboratory setting. *Neuropsychobiology*, 28(1–2), 76–81.
- Kokkonen, M., & Pulkkinen, L. (2001). Extraversion and neuroticism as antecedents of emotion regulation and dysregulation in adulthood. *European Journal of Personality*, 15(6), 407–424.
- Kraus, S., & Sears, S. (2009). Measuring the Immeasurables: development and initial validation of the Self-Other Four Immeasurables (SOFI) scale based on Buddhist teachings on loving kindness, compassion, joy, and equanimity. *Social Indicators Research*, 92(1), 169–181.
- Kross, E., & Ayduk, Ö. (2008). Facilitating adaptive emotional analysis: distinguishing distanced-analysis of depressive experiences from immersed-analysis and distraction. *Personality & Social Psychology Bulletin*, 34(7), 924–938.
- Kross, E., Ayduk, Ö., & Mischel, W. (2005). When asking “why” does not hurt. Distinguishing rumination from reflective processing of negative emotions. *Psychological Science*, 16(9), 709–715.
- Lazarus, R. S. (1991). *Emotion and adaptation*. New York: Oxford University Press.
- Ledoux, J. (1996). *The emotional brain: the mysterious underpinnings of emotional life*. New York, NY: Simon & Schuster.
- Levenson, R. W., Ekman, P., & Ricard, M. (2012). Meditation and the startle response: a case study. *Emotion*, 12(3), 650–658.
- Lévesque, J., Eugene, F., Joannette, Y., Paquette, V., Mensour, B., Beaudoin, G., et al. (2003). Neural circuitry underlying voluntary suppression of sadness. *Biological Psychiatry*, 53(6), 502–510.
- Lundman, B., Strandberg, G., Eisemann, M., Gustafson, Y., & Brulin, C. (2007). Psychometric properties of the Swedish version of the Resilience Scale. *Scandinavian Journal of Caring Sciences*, 21(2), 229–237.
- Lutz, A., Dunne, J. D., & Davidson, R. J. (2007). Meditation and the neuroscience of consciousness: an introduction. In P. D. Zelazo, M. Moscovitch, & E. Thompson (Eds.), *The Cambridge handbook of consciousness* (1st ed., pp. 499–551). Cambridge, UK: Cambridge University Press.
- Lutz, A., Brefczynski-Lewis, J., Johnstone, T., & Davidson, R. J. (2008). Regulation of the neural circuitry of emotion by compassion meditation: effects of meditative expertise. *PLoS One*, 3(3), e1897.
- Mack, J. W., Nilsson, M., Balboni, T., Friedlander, R. J., Block, S. D., Trice, E., & Prigerson, H. G. (2008). Peace, Equanimity, and Acceptance in the Cancer Experience (PEACE): validation of a scale to assess acceptance and struggle with terminal illness. *Cancer*, 112(11), 2509–2517.
- Makransky, J. (2007). *Awakening through love: unveiling your deepest goodness*. Somerville: Wisdom Publications.
- Mason, O., & Hargreaves, I. (2001). A qualitative study of mindfulness-based cognitive therapy for depression. *British Journal of Medical Psychology*, 74, 197–212.
- McEwen, B. S. (1998). Protective and damaging effects of stress mediators. *The New England Journal of Medicine*, 338(3), 171–179.
- McEwen, B. S. (2008). Central effects of stress hormones in health and disease: understanding the protective and damaging effects of stress and stress mediators. *European Journal of Pharmacology*, 583(2–3), 174–185.
- McEwen, B. S., & Seeman, T. E. (1999). Protective and damaging effects of mediators of stress. Elaborating and testing the concepts of allostasis and allostatic load. *Annals of the New York Academy of Sciences*, 896, 30–47.
- McRae, K., Ochsner, K. N., & Gross, J. J. (2010). The reason in passion: a social cognitive neuroscience approach to emotion regulation. In K. D. Vohs & R. F. Baumeister (Eds.), *Handbook of self-regulation, second edition: research, theory, and applications* (pp. 186–203). New York: The Guilford Press.
- Moore, A., & Malinowski, P. (2009). Meditation, mindfulness and cognitive flexibility. *Consciousness and Cognition*, 18(1), 176–186.
- Najström, M., & Jansson, B. (2007). Skin conductance responses as predictor of emotional responses to stressful life events. *Behaviour Research and Therapy*, 45(10), 2456–2463.
- Nanamoli, B. (1991). *The path of purification*. Onalaska: Buddhist Publication Society Pariyatti Editions.
- Northoff, G., & Berman, F. (2004). Cortical midline structures and the self. *Trends in Cognitive Sciences*, 8(3), 102–107.
- Northoff, G., Heinzel, A., de Greck, M., Berman, F., Dobrowolny, H., & Panksepp, J. (2006). Self-referential processing in our brain—a meta-analysis of imaging studies on the self. *NeuroImage*, 31(1), 440–457.
- Nyanaponika Thera (1962). *The heart of Buddhist meditation: a handbook of mental training based on the Buddha's way of mindfulness*. London: Rider.
- Ochsner, K. N., & Gross, J. J. (2005). The cognitive control of emotion. *Trends in Cognitive Sciences*, 9(5), 242–249.
- Ochsner, K. N., Bunge, S. A., Gross, J. J., & Gabrieli, J. D. E. (2002). Rethinking feelings: an fMRI study of the cognitive regulation of emotion. *Journal of Cognitive Neuroscience*, 14(8), 1215–1229.
- Ochsner, K. N., Ray, R. D., Cooper, J. C., Robertson, E. R., Chopra, S., Gabrieli, J. D. E., et al. (2004). For better or for worse: neural systems supporting the cognitive down- and up-regulation of negative emotion. *NeuroImage*, 23(2), 483–499.
- Ohira, H., Nomura, M., Ichikawa, N., Isowa, T., Iidaka, T., Sato, A., et al. (2006). Association of neural and physiological responses during voluntary emotion suppression. *NeuroImage*, 29(3), 721–733.
- Olendzki, A. (2011). The construction of mindfulness. *Contemporary Buddhism*, 12(1), 55–70.
- Ortner, C. N., Kilner, S. J., & Zelazo, P. D. (2007). Mindfulness meditation and reduced emotional interference on a cognitive task. *Motivation and Emotion*, 31(4), 271–283.
- Pace, T. W. W., Mletzko, T. C., Alagbe, O., Musselman, D. L., Nemeroff, C. B., Miller, A. H., et al. (2006). Increased stress-induced inflammatory responses in male patients with major depression and increased early life stress. *American Journal of Psychiatry*, 163(9), 1630–1633.
- Pace, T. W. W., Negi, L. T., Adame, D. D., Cole, S. P., Sivilli, T. I., Brown, T. D., et al. (2009). Effect of compassion meditation on neuroendocrine, innate immune and behavioral responses to psychosocial stress. *Psychoneuroendocrinology*, 34(1), 87–98.
- Pagnoni, G., Cekic, M., & Guo, Y. (2008). “Thinking about nothing”: neural correlates of conceptual processing during Zen meditation. *PLoS One*, 3(9), e3083.
- Parkinson, B., & Totterdell, P. (1999). Classifying affect-regulation strategies. *Cognition and Emotion*, 13(3), 277–303.
- Perlman, D. M., Salomons, T. V., Davidson, R. J., & Lutz, A. (2010). Differential effects on pain intensity and unpleasantness of two meditation practices. *Emotion*, 10(1), 65–71.
- Porges, S. W. (2007). The polyvagal perspective. *Biological Psychology*, 74(2), 116–143.
- Porges, S. W. (2011). *The polyvagal theory: neurophysiological foundations of emotions, attachment, communication, and self-regulation (Norton Series on Interpersonal Neurobiology)*. New York: Norton.
- Pu, J., Schmeichel, B. J., & Demaree, H. A. (2010). Cardiac vagal control predicts spontaneous regulation of negative emotional expression and subsequent cognitive performance. *Biological Psychology*, 84(3), 531–540.
- Purdon, C. (1999). Thought suppression and psychopathology. *Behaviour Research and Therapy*, 37(11), 1029–1054.

- Robinson, J. L., & Demaree, H. A. (2009). Experiencing and regulating sadness: physiological and cognitive effects. *Brain and Cognition*, *70*(1), 13–20.
- Ruiz-Padial, E., Vila, J., & Thayer, J. F. (2011). The effect of conscious and non-conscious presentation of biologically relevant emotion pictures on emotion modulated startle and phasic heart rate. *International Journal of Psychophysiology*, *79*(3), 341–346.
- Safran, J., & Segal, Z. V. (1990). *Interpersonal process in cognitive therapy*. New York: Basic Books.
- Sahdra, B. K., Shaver, P. R., & Brown, K. W. (2010). A scale to measure nonattachment: a Buddhist complement to Western research on attachment and adaptive functioning. *Journal of Personality Assessment*, *92*(2), 116–127.
- Salzberg, S. (1995). *Loving-kindness: the revolutionary art of happiness*. Boston: Shambhala Publications.
- Schaefer, S. M., Jackson, D. C., Davidson, R. J., Aguirre, G. K., Kimberg, D. Y., & Thompson-Schill, S. L. (2002). Modulation of amygdalar activity by the conscious regulation of negative emotion. *Journal of Cognitive Neuroscience*, *14*(6), 913–921.
- Schooler, J. W., Smallwood, J., Christoff, K., Handy, T. C., Reichle, E. D., & Sayette, M. A. (2011). Meta-awareness, perceptual decoupling and the wandering mind. *Trends in Cognitive Sciences*, *15*(7), 319–326.
- Schuyler, B. S., Kral, T. R. A., Jacquart, J., Burghy, C. A., Weng, H. Y., Perlman, D. M., et al. (2012). Temporal dynamics of emotional responding: amygdala recovery predicts emotional traits. *Social Cognitive and Affective Neuroscience*. doi:10.1093/scan/nss131.
- Segal, Z. V., Williams, J. M. G., & Teasdale, J. D. (2001). *Mindfulness-based cognitive therapy for depression: a new approach to preventing relapse*. New York: The Guilford Press.
- Shapiro, S. L., Carlson, L. E., Astin, J. A., & Freedman, B. (2006). Mechanisms of mindfulness. *Journal of Clinical Psychology*, *62*(3), 373–386.
- Sheppes, G., Catran, E., & Meiran, N. (2009). Reappraisal (but not distraction) is going to make you sweat: physiological evidence for self-control effort. *International Journal of Psychophysiology*, *71*(2), 91–96.
- Slagter, H. A., Davidson, R. J., & Lutz, A. (2011). Mental training as a tool in the neuroscientific study of brain and cognitive plasticity. *Frontiers in Human Neuroscience*, *5*, 17.
- Smith, T. W., Cribbet, M. R., Nealey-Moore, J. B., Uchino, B. N., Williams, P. G., Mackenzie, J., et al. (2011). Matters of the variable heart: respiratory sinus arrhythmia response to marital interaction and associations with marital quality. *Journal of Personality and Social Psychology*, *100*(1), 103–119.
- Stellar, J. E., Manzo, V. M., Kraus, M. W., & Keltner, D. (2012). Class and compassion: socioeconomic factors predict responses to suffering. *Emotion*, *12*(3), 449–459.
- Steptoe, A., Hamer, M., & Chida, Y. (2007). The effects of acute psychological stress on circulating inflammatory factors in humans: a review and meta-analysis. *Brain, Behavior, and Immunity*, *21*(7), 901–912.
- Sterling, P., & Eyer, J. (1988). Allostasis: a new paradigm to explain arousal pathology. In S. Fisher & J. Reason (Eds.), *Handbook of life stress, cognition and health* (pp. 629–649). New York: Wiley.
- Sternberg, E. (2000). *The balance within: the science connecting health and emotions*. New York: Freeman.
- Suls, J., & Martin, R. (2005). The daily life of the garden-variety neurotic: reactivity, stressor exposure, mood spillover, and maladaptive coping. *Journal of Personality*, *73*(6), 1485–1509.
- Taylor, V. A., Grant, J. A., Daneault, V., Scavone, G., Breton, E., Roffe-Vidal, S., et al. (2011). Impact of mindfulness on the neural responses to emotional pictures in experienced and beginner meditators. *NeuroImage*, *57*(4), 1524–1533.
- Teasdale, J. D. (1999). Metacognition, mindfulness and the modification of mood disorders. *Clinical Psychology & Psychotherapy*, *6*(2), 146–155.
- Teasdale, J. D., Segal, Z. V., & Williams, J. M. (1995). How does cognitive therapy prevent depressive relapse and why should attentional control (mindfulness) training help? *Behaviour Research and Therapy*, *33*(1), 25–39.
- Teasdale, J. D., Moore, R. G., Hayhurst, H., Pope, M., Williams, S., & Segal, Z. V. (2002). Metacognitive awareness and prevention of relapse in depression: empirical evidence. *Journal of Consulting and Clinical Psychology*, *70*(2), 275–287.
- Thanissaro Bhikkhu (1996). *Wings to awakening: an anthology from the Pali Canon*. Barre: Dhamma Dana Publications.
- Thayer, J. F., & Lane, R. D. (2000). A model of neurovisceral integration in emotion regulation and dysregulation. *Journal of Affective Disorders*, *61*(3), 201–216.
- Thayer, J. F., Åhs, F., Fredrikson, M., Sollers, J. J., III, & Wager, T. D. (2012). A meta-analysis of heart rate variability and neuroimaging studies: implications for heart rate variability as a marker of stress and health. *Neuroscience & Biobehavioral Reviews*, *36*(2), 747–756.
- Thrangun Rinpoche, K. (2002). *Essential practice*. Ithaca: Snow Lion Publications.
- Travis, F., & Shear, J. (2010). Focused attention, open monitoring and automatic self-transcending: categories to organize meditations from Vedic, Buddhist and Chinese traditions. *Consciousness and Cognition*, *19*(4), 1110–1118.
- Tsering, G. T. (2006). *Buddhist psychology: The foundation of Buddhist thought, vol. 3*. Somerville: Wisdom Publications.
- Vago, D. R., & Nakamura, Y. (2011). Selective attentional bias towards pain-related threat in fibromyalgia: preliminary evidence for effects of mindfulness meditation training. *Cognitive Therapy and Research*, *35*(6), 581–594.
- Van Tongeren, D. R., & Green, J. D. (2010). Combating meaninglessness: on the automatic defense of meaning. *Personality & Social Psychology Bulletin*, *36*(10), 1372–1384.
- Varela, F. J., Thompson, E. T., & Rosch, E. (1992). *The embodied mind: cognitive science and human experience*. Cambridge: The MIT Press.
- Vohs, K. D., & Baumeister, R. F. (2010). *Handbook of self-regulation, second edition: research, theory, and applications*. New York: The Guilford Press.
- Volkhov, R. N., & Demaree, H. A. (2010). Spontaneous emotion regulation to positive and negative stimuli. *Brain and Cognition*, *73*(1), 1–6.
- von Känel, R., Kudielka, B. M., Preckel, D., Hanebuth, D., & Fischer, J. E. (2006). Delayed response and lack of habituation in plasma interleukin-6 to acute mental stress in men. *Brain, Behavior, and Immunity*, *20*(1), 40–48.
- Vuilleumier, P. (2005). How brains beware: neural mechanisms of emotional attention. *Trends in Cognitive Sciences*, *9*(12), 585–594.
- Wallace, B. A. (2006). *The attention revolution: unlocking the power of the focused mind*. Somerville: Wisdom Publications.
- Wallace, B. A. (2010). *The four immeasurables: practices to open the heart*. Ithaca: Snow Lion Publications.
- Wallace, B. A. (2011). *Minding closely: the four applications of mindfulness*. Ithaca: Snow Lion Publications.
- Whalen, P. J. (1998). Fear, vigilance, and ambiguity: initial neuroimaging studies of the human amygdala. *Current Directions in Psychological Science*, *7*(6), 177–188.
- Whalen, P. J., & Phelps, E. A. (Eds.). (2009). *The human amygdala*. New York: The Guilford Press.
- Williams, J. M. G. (2010). Mindfulness and psychological process. *Emotion*, *10*(1), 1–7.
- Williams, J. M. G., & Kabat-Zinn, J. (2011). Mindfulness: diverse perspectives on its meaning, origins, and multiple applications at the intersection of science and Dharma. *Contemporary Buddhism*, *12*(1), 1–18.
- Wirtz, P. H., von Känel, R., Emini, L., Suter, T., Fontana, A., & Ehler, U. (2007). Variations in anticipatory cognitive stress appraisal and differential proinflammatory cytokine expression in response to acute stress. *Brain, Behavior, and Immunity*, *21*(6), 851–859.
- Yamakawa, K., Matsunaga, M., Isowa, T., Kimura, K., Kasugai, K., Yoneda, M., et al. (2009). Transient responses of inflammatory cytokines in acute stress. *Biological Psychology*, *82*(1), 25–32.